

UC Irvine

UC Irvine Previously Published Works

Title

Effects of febrile seizures in the immature rat model on interneuronal parvalbumin (PV) expression in the dentate gyrus

Permalink

<https://escholarship.org/uc/item/2091n2vn>

Journal

EPILEPSIA, 40

ISSN

0013-9580

Authors

Bender, RA
Baram, TZ

Publication Date

1999

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

2.144

EFFECTS OF FEBRILE SEIZURES IN THE IMMATURE RAT MODEL ON INTERNEURONAL PARVALBUMIN (PV) EXPRESSION IN THE DENTATE GYRUS

Roland A Bender, University of California, Irvine, IRVINE, CA; Tallie Z Baram, UCI, Irvine, CA

RATIONALE: A febrile seizure model, recently established in the immature rat (Baram et al., Brain Res, 1997) may serve as a valuable tool to study the effects of these seizures on the developing hippocampus. Loss of dentate gyrus PV-immunoreactivity after experimentally induced seizures has been reported in other models, and has been interpreted as either loss of interneurons or their dysfunction, leading to granule cell disinhibition. **OBJECTIVE:** To test the hypothesis that PV expression is persistently down-regulated after febrile seizures. **METHODS:** Hyperthermic seizures (mean duration 21 min) were induced on postnatal day 10. PV-mRNA and protein levels were evaluated using in situ hybridization and immunocytochemistry (ICC), respectively. Quantitative analysis of PV-expressing neurons in dentate gyrus was achieved blindly, using stereologic approaches. **RESULTS:** The

number of PV expressing neurons of experimental and control groups were not significantly different at 4 hours, 1 week or 4 weeks after seizures. For example, at 1 week, PV-immunoreactive neurons totaled 4353 ± 298 and 4361 ± 471 in controls and experimentals, respectively ($n = 3$); corresponding values at 4 weeks were 6362 ± 611 and 6378 ± 734 ($n = 4$). In addition, PV-mRNA expression (at 4 hours and 4 weeks) was not down-regulated after the hyperthermic seizures. **CONCLUSION:** In this model of developmental seizures, down-regulation of PV expression is not observed, consistent with (compensatory?) increased inter-neuron-mediated inhibition (Chen et al., Soc. Neurosci. abst., 1998) reported in the same model. The molecular mechanisms of the long-lasting enhanced susceptibility of rats experiencing febrile seizures to hippocampal convulsants (Dube et al., this meeting) require further study. (Supported by NIH NS35439)